Upper Gastrointestinal Bleeding From Aberrant Right Subclavian Artery-Esophageal Fistula

Shira Peress, MD,¹ Wa'el Tuqan, MD,² Austin Thomas, MD²

¹The University of Queensland Faculty of Medicine, Ochsner Clinical School, New Orleans, LA ²Department of Gastroenterology, Ochsner Clinic Foundation, New Orleans, LA

Background: The most common aortic arch abnormality is an aberrant right subclavian artery (ARSA). ARSA-esophageal fistula is a rare sequela that can present with a life-threatening upper gastrointestinal (GI) bleed.

Case Report: We report the case of an 88-year-old male who presented with signs of upper GI bleeding. Esophagogastroduodenoscopy demonstrated extrinsic compression of the upper third of the esophagus with ulceration. Imaging studies revealed ARSA posterior to the esophagus with pseudoaneurysm formation. These findings confirmed an upper GI bleed secondary to ARSA-esophageal fistula. The patient underwent prompt embolization of the ARSA pseudoaneurysm, followed a few days later by coil embolization of the ARSA pseudoaneurysm. Despite these interventions, the patient continued to have bleeding with anemia. He and his family opted to avoid any further interventions and instead pursued comfort care. The patient was discharged to hospice and died 3 months later.

Conclusion: ARSA-esophageal fistula is a rare but potentially lethal cause of upper GI bleeding. Initial signs and symptoms can be subtle, but the presence of a GI bleed requires immediate stabilization. Surgical interventions have been shown to have longer-lasting success, but endovascular repair may be an option for patients who are deemed unfit for surgery but still require prompt stabilization. Regardless of the intervention, mortality rates for ARSA-esophageal fistula are high.

Keywords: Aberrant right subclavian artery, aneurysm, esophageal fistula, gastrointestinal hemorrhage

Address correspondence to Austin Thomas, MD, Department of Gastroenterology, Ochsner Clinic Foundation, 1514 Jefferson Hwy., New Orleans, LA 70121. Tel: (504) 842-4015. Email: aauthomas@ochsner.org

INTRODUCTION

Aberrant right subclavian artery (ARSA), or arteria lusoria, is the most common aortic arch abnormality, with a prevalence of 0.2% to 2.5%.^{1,2} ARSA is typically an incidental finding that seldom causes any symptoms.³ However, ARSA is prone to aneurysmal development that can remain asymptomatic or cause compressive symptoms, including shortness of breath and dysphagia lusoria secondary to extrinsic compression of the esophagus.³ The recurrent laryngeal nerve is also prone to a degree of constriction from ARSA, causing progressive hoarseness. An even rarer consequence is the development of ARSA-esophageal fistula that can present as a life-threatening upper gastrointestinal (GI) bleed, so prompt diagnosis and management are essential for patient survival.

CASE REPORT

An 88-year-old male presented to the emergency department (ED) with a 1-day history of shortness of breath. He reported one episode of dark black, bloody stools 2 days prior. He also reported dysphagia for 6 months and hoarseness of voice for 1 week. He had an extensive cardiovascular history, including aortic stenosis status post transcatheter

aortic valve replacement, thoracic aortic aneurysm (TAA), atrial fibrillation on warfarin, and congestive heart failure. A month and a half prior to his presentation, computed tomography (CT) angiography of the chest to evaluate the TAA demonstrated an ascending TAA and saccular pseudoaneurysm formation from an ARSA, resulting in marked mass effect and displacement of the esophagus (Figure 1). Bilateral carotid subclavian bypass was recommended; however, the patient declined surgery at the time.

On presentation to the ED, he was hemodynamically stable. Laboratory workup revealed hemoglobin of 6.5 g/dL from a baseline of 11.0 g/dL. Prothrombin time was 17.1 seconds, with an international normalized ratio (INR) of 1.8. Following initial resuscitation with 2 units of packed red blood cells and intravenous pantoprazole, esophagogastroduodenoscopy (EGD) was performed and demonstrated extrinsic compression of the upper third of the esophagus, with associated areas of ulceration and visible underlying vasculature (Figure 2). EGD findings were concerning for development of ARSA-esophageal fistula and the potential for subsequent bleeding. Vascular surgery performed urgent embolization of the ARSA using Amplatzer plugs.

Figure 1. Computed tomography angiography of the chest demonstrates an ascending thoracic aortic aneurysm with saccular pseudoaneurysm formation. Arrows point to the aneurysm compressing the esophagus.

Figure 2. Esophagogastroduodenoscopy demonstrates (A) external compression into the lumen of the esophagus and (B) the inferior part of the compression with ulceration (arrows) and possible visible vessel.

Figure 3. Interventional radiology imaging demonstrates (A) precoiled pseudoaneurysm and (B) postcoiled pseudoaneurysm.

The patient initially did well and had no further signs of bleeding. He was discharged 2 days after the procedure with hemoglobin of 7.7 g/dL and INR of 1.2. One day after discharge, he presented again to the ED with worsening fatigue, weakness, and shortness of breath. He had 2 additional episodes of melena and a drop in hemoglobin to 6.5 g/dL. He was readmitted, and repeat CT angiography demonstrated a new bleed within the pseudoaneurysm sac. In conjunction with vascular surgery, interventional radiology performed a coil embolization of the ARSA pseudoaneurysm (Figure 3).

Despite these interventions, the patient continued to have ongoing signs of bleeding with anemia. He required multiple blood transfusions for stabilization, ultimately receiving

Figure 4. Schematic representation of aberrant right subclavian artery. (Image courtesy of Dr Vincent Tatco, Radiopaedia.org, rID: 52193 radiopaedia.org/cases/development-of-aberrant-right-subclavian-artery-illustration)

5 units of packed red blood cells that helped improve his hemoglobin to 9.2 g/dL and hematocrit to 28.5%. After multidisciplinary team discussions, the patient and his family opted to avoid any further workup or surgical interventions and instead pursued comfort care. He was discharged to hospice 4 days after being readmitted and died 3 months later.

DISCUSSION

When ARSA is present, the right subclavian artery branches distal to the left subclavian artery and serves as the fourth branch of the left-sided aortic arch (Figure 4). From there, the artery travels posterior to the esophagus in 80% of cases but can also be found between the trachea and esophagus in 15% of cases and anterior to the trachea in 5% of cases.⁴ ARSA has been reported to have a female predominance as high as 3:1, although other studies suggest no significant sex discrepancy.⁵ ARSA can be associated with other medical conditions and is seen in up to 25% of patients with esophageal atresia and in 16% to 39% of patients with Down syndrome.^{6,7}

Despite its prevalence in the general population, ARSA is frequently asymptomatic and discovered incidentally on imaging. However, ARSA can present with dysphagia secondary to esophageal compression, a phenomenon known as dysphagia lusoria.⁸ Depending on the exact location, ARSA can cause some degree of compression of the recurrent laryngeal nerve, resulting in hoarseness, a condition known as Ortner syndrome.⁹ ARSA can also cause compression of the trachea, resulting in shortness of breath. With advancing age, ARSA is prone to aneurysmal development or sclerosis.³ The late presentation of dysphagia in our elderly patient is not uncommon, as dysphagia is more frequently seen in the elderly compared to younger patients with ARSA and can be attributed to decreased flexibility of the esophagus or increasing compression from the developing aneurysm. Some degree of atherosclerosis contributing to rigidity of the ARSA is also possible.¹⁰

In our review of literature, we found 35 reported cases of ARSA-esophageal fistula.^{1,11-43} Of the 35 cases, 33 cases are included in our summary table, while 2 cases were excluded due to inability to access the primary source or the source being in a language other than English (Table).^{13,20} Six cases were in the pediatric population, with patient ages ranging from 5 months to 11 years.^{22,28,30,34,37} Although diagnosis is typically made with CT imaging, endoscopic visualization may be the initial diagnostic test to exclude other causes, particularly in patients who are hemodynamically stable at the time of presentation. The most common etiology for development of ARSA-esophageal fistula appeared to be secondary to compression, friction, or pressure necrosis from recent instrumentation with an endotracheal tube, nasogastric tube, or tracheostomy tube, seen in 14 of the 33 summarized cases.^{14,15,17,19,21,22,24,27,29,31,38-41} Fistula development from prior placement of esophageal stent was noted in 5 cases.^{34,36,42,43} Other reported causes for fistula formation included placement of a salivary bypass tube and a possible consequence of gastric pull-up surgery.^{1,26}

Fistula formation attributable to ARSA aneurysm with resultant GI bleeding is rare. Only 10 other cases attributed to this etiology have been reported.^{11,12,16,18,19,23,25,33,35,42} Only 1 patient with upper GI bleed secondary to ARSA aneurysm survived all bleeding episodes, and the patient required balloon tamponade with surgery to replace the aorta.³³ Our patient survived the initial bleeding event, but died 3 months later, as he declined definitive surgical intervention to stop the recurrent bleeding.

Bleeding from ARSA-esophageal fistula is life-threatening, with a high mortality rate at initial presentation, regardless of etiology. Of the 33 summarized cases, thirteen cases survived all episodes of bleeding (39%).^{21,22,24,27,28,30,31,33,34,36-38,43} The severity of bleeding, in the setting of several comorbid conditions, likely contributed to the high mortality rate.

Opinions conflict regarding acute treatment for bleeding from ARSA-esophageal fistula secondary to aneurysm, with the historic approach being excision with reconstruction of the aorta and esophagus.³³ A more recent approach (2000) is placement of an endovascular stent, but success with this approach is questionable because of the rates of infection that may result in the need for further surgery.⁴⁴ Reported cases in adults involved treatment with surgery alone in 5 cases,^{11,16,26,30,38} endovascular repair alone in 2 cases,^{41,42} and a combination of endovascular and surgical intervention in 6 cases.^{28,29,31,35,36,43} Balloon tamponade was performed in 16 cases,^{1,14,15,21,22,24,25,27-29,31-33,35-37} with 3 of those 16 performed solely with endovascular repair.^{1,27,32} Six of the 11 patients who underwent either surgery alone or surgery with endovascular repair survived (55%),^{28,30,31,36,38,43} whereas only 1 of the 5 cases of endovascular repair with or without balloon tamponade survived (20%).²⁷ Although the higher percentage of survival among patients who had surgery with or without endovascular repair compared to the 5 patients who underwent endovascular repair without surgery might suggest a better outcome with surgery, this discrepancy is likely confounded by patient factors and comorbidities for which surgical intervention is usually deferred, as well as the relatively low number of patient cases reported to have been treated with endovascular repair without surgery.

Table. Summary of Reported Cases of Aberrant Right Subclavian Artery-Esophageal Fistula

Study	Age, Sex	Medical History	Etiology	Treatment	Outcome
Lynn, 1969 ¹¹	57, M	Atherosclerosis, hypertension, angina	ARSA aneurysm	Surgery	Fatal
Reynes et al, 1976 ¹²	72, F	Mediastinal mass	ARSA aneurysm	None	Fatal
Livesay et al, 1982 ¹⁴	25, M	Motor vehicle accident with traumatic head injury	Endotracheal tube, nasogastric tube, and tracheostomy	Balloon tamponade, surgery	Fatal
Belkin et al, 1984 ¹⁵	27, M	Head and neck squamous cell carcinoma	Nasogastric tube	Balloon tamponade, surgery	Fatal
Edwards et al, 1984 ¹⁶	81, M	Mass in right upper hemithorax (tortuous innominate artery)	ARSA aneurysm	Surgery	Fatal
Gossot et al, 1985 ¹⁷	72, F	Type I aortic dissection with endoluminal prosthesis, giant cell arteritis	Endotracheal tube, nasogastric tube, and tracheostomy	None	Fatal
Kullnig, 1989 ¹⁸	66, M	Hypertension	ARSA aneurysm	None	Fatal
Stone et al, 1990 ¹⁹	72, M	Smoker, status post cardiopulmonary bypass surgery	ARSA aneurysm, tracheostomy	None	Fatal
Hirakata et al, 1991 ²¹	55, M	Status post subtotal esophagectomy for esophageal cancer	Nasogastric tube, radiation arteritis, and other surgical trauma	Balloon tamponade	Survived
Miller et al, 1996 ²²	11, F	Status post craniotomy for intracerebral hemorrhage	Endotracheal tube, nasogastric tube	Balloon tamponade, surgery	Survived
Singha et al, 1998 ²³	82, M	N/R	ARSA aneurysm	None	Fatal
Feugier et al, 2002 ²⁴	24, M	Polytrauma, burns, alcohol abuse	Nasogastric tube and tracheotomy	Balloon tamponade, surgery	Survived
Lehmann et al, 2006 ²⁵	78, M	N/R	ARSA aneurysm	Balloon tamponade	Fatal
Millar et al, 2007 ²⁶	57, M	Status post esophagectomy with gastric pull-up for esophageal cancer	Pressure from gastric pull-up vs foreign body	Surgery	Fatal (survived initial hemorrhage but died 18 days later from additional bleed)
Inman et al, 2008 ¹	63, M	Supraglottic squamous cell carcinoma	Salivary bypass tube	Endovascular repair, balloon tamponade	Fatal
Magagna et al, 2008 ²⁷	73, F	Status post laryngectomy and tracheostomy for laryngeal carcinoma	Tracheostomy	Endovascular repair, balloon tamponade	Survived
Fuentes et al, 2010 ²⁸	3, F	Esophageal atresia type III	Esophageal prosthesis	Endovascular repair, balloon tamponade, surgery	Survived
Chapman et al, 2010 ²⁹	34, F	Motor vehicle accident with trauma	Endotracheal tube, nasogastric tube, and tracheostomy	Endovascular repair, balloon tamponade, surgery	Fatal
Situma et al, 2011 ³⁰	5 months, F	Esophageal atresia with distal fistula	Status post colonic esophageal grafting	Surgery	Survived
Jain et al, 2012 ³¹	57, F	Status post cardiopulmonary bypass surgery for scimitar syndrome	Endotracheal tube, nasogastric tube, and tracheostomy	Endovascular repair, balloon tamponade, surgery	Survived

Table. Continued

Study	Age, Sex	Medical History	Etiology	Treatment	Outcome
Pop et al, 2012 ³²	67, M	Status post transhiatal esophagectomy for esophageal cancer, status post pneumonectomy for lung cancer, neuroendocrine liver cancer	Damage to arterial wall from cervical sepsis and/or staples from gastric tubing	Endovascular repair, balloon tamponade	Fatal
Takahashi et al, 2013 ³³	63, M	N/R	Infected ARSA aneurysm	Balloon tamponade, surgery	Survived
Lo et al, 2013 ³⁴	16 months, N/R	Esophageal atresia with distal fistula	Esophageal and Polyflex airway stent	Endovascular repair, Bougienage tamponade	Fatal
	18 months, N/R	Esophageal atresia, duodenal atresia	Esophageal stent	Surgery, hydrostatic dilator	Survived
Morisaki et al, 2014 ³⁵	74, F	Rheumatic arthritis	ARSA aneurysm	Endovascular repair, balloon tamponade, surgery	Fatal
Hosn et al, 2014 ³⁶	29, F	Sleeve gastrectomy	Esophageal stent	Endovascular repair, balloon tamponade, surgery	Survived
Joynt and Grifka, 2015 ³⁷	17 months, F	N/R	Spontaneous development of fistula	Balloon tamponade, surgery	Survived
Watanabe et al, 2015 ³⁹	55, M	Intracranial hemorrhage	Nasogastric tube and tracheostomy	None	Fatal
Oliveira et al, 2016 ³⁸	20, M	Motor vehicle accident with polytrauma	Endotracheal tube and nasogastric tube	Surgery	Survived
Kudose et al, 2017 ⁴⁰	20, F	VATER (vertebral, anal, tracheal, esophageal, and renal) association, primary pulmonary hypertension, diabetes mellitus, atrial septal defect, status post 3 lung transplants	Tracheostomy	None	Fatal
Shires and Rohrer, 2018 ⁴¹	44, F	Gastroesophageal reflux disease, hypertension, pneumonia	Nasogastric tube, endotracheal tube, and tracheostomy	Endovascular repair	Fatal
Zheng et al, 2019 ⁴²	67, M	Esophageal and laryngeal cancer, hypertension	Esophageal stent and pseudoaneurysm	Endovascular repair	Fatal
Merlo et al, 2020 ⁴³	29, F	Tracheoesophageal fistula, ventriculoperitoneal shunt for hydrocephalus	Esophageal stent	Endovascular repair, surgery	Survived
Present case, 2021	88, M	Hypertension, transcatheter aortic valve replacement, thoracic aortic aneurysm	ARSA aneurysm	Endovascular repair	Fatal (survived initial event, but died 3 months after discharge from recurrent bleeds)

Note: Age is given in years unless otherwise indicated.

ARSA, aberrant right subclavian artery; F, female; M, male; N/R, not reported.

CONCLUSION

ARSA-esophageal fistula is an uncommon and potentially fatal cause of GI bleeding. Patients with known ARSA aneurysm should elicit a high index of suspicion, particularly those who present with concomitant worsening dysphagia. Given the high mortality with active bleeding, endovascular repair is a potential alternative for patients who are not suitable for surgery because of their comorbidities or high surgical risk. However, even with intervention, mortality remains high.

ACKNOWLEDGMENTS

The authors have no financial or proprietary interest in the subject matter of this article.

REFERENCES

- Inman JC, Kim P, McHugh R. Retroesophageal subclavian artery-esophageal fistula: a rare complication of a salivary bypass tube. *Head Neck*. 2008;30(8):1120-1123. doi: 10.1002/hed.20854
- Nakatani T, Tanaka S, Mizukami S, Okamoto K, Shiraishi Y, Nakamura T. Retroesophageal right subclavian artery originating from the aortic arch distal and dorsal to the left subclavian artery. *Ann Anat*. 1996;178(3):269-271. doi: 10.1016/S0940-9602(96)80064-1
- Abraham V, Mathew A, Cherian V, Chandran S, Mathew G. Aberrant subclavian artery: anatomical curiosity or clinical entity. *Int J Surg*. 2009;7(2):106-109. doi: 10.1016/j.ijsu.2009.01.009
- Gomes MM, Bernatz PE, Forth RJ. Arteriosclerotic aneurysm of an aberrant right subclavian artery. *Dis Chest*. 1968;54(6):549-552. doi: 10.1378/chest.54.6.549
- Molz G, Burri B. Aberrant subclavian artery (arteria lusoria): sex differences in the prevalence of various forms of the malformation. Evaluation of 1378 observations. *Virchows Arch A Pathol Anat Histol*. 1978;380(4):303-315. doi: 10.1007/BF00431315
- Berthet S, Tenisch E, Miron MC, et al. Vascular anomalies associated with esophageal atresia and tracheoesophageal fistula. *J Pediatr*. 2015;166(5):1140-1144.e2. doi: 10.1016/j.jpeds.2015.01.038
- Chaoui R, Heling KS, Sarioglu N, Schwabe M, Dankof A, Bollmann R. Aberrant right subclavian artery as a new cardiac sign in second- and third-trimester fetuses with Down syndrome. *Am J Obstet Gynecol*. 2005;192(1):257-263. doi: 10.1016/j.ajog.2004.06.080
- Levitt B, Richter JE. Dysphagia lusoria: a comprehensive review. *Dis Esophagus*. 2007;20(6):455-460. doi: 10.1111/j.1442-2050.2007.00787.x
- Bickle IC, Kelly BE, Brooker DS. Ortner's syndrome: a radiological diagnosis. *Ulster Med J*. 2002;71(1):55-56.
- Kantarceken B, Bulbuloglu E, Yuksel M, Cetinkaya A. Dysphagia lusoria in elderly: a case report. *World J Gastroenterol*. 2004;10(16):2459-2460. doi: 10.3748/wjg.v10.i16.2459
- Lynn RB. Kommerell's diverticulum with esophago-arterial fistula. *Can J Surg*. 1969;12(3):331-333.
- Reynes JB, Errasti CA, Ercoreca FJB, Oñate Landa A. Letter: aneurysm of aberrant right subclavian artery with esophageal perforation. *Chest*. 1976;70(1):105. doi: 10.1378/chest.70.1.105a
- Merchant FJ, Nichols RL, Bombeck CT. Unusual complication of nasogastric esophageal intubation-erosion into an aberrant right subclavian artery. *J Cardiovasc Surg (Torino)*. 1977;18(2):147-150.
- Livesay JJ, Michals AA, Dainko EC. Anomalous right subclavian arterial esophageal fistula: an unusual complication of tracheostomy. *Tex Heart Inst J*. 1982;9(1):105-108.
- Belkin RI, Keller FS, Everts EC, Röscher J. Aberrant right subclavian artery-esophageal fistula: a cause of overwhelming upper gastrointestinal hemorrhage. *Cardiovasc Intervent Radiol*. 1984;7(2):87-89. doi: 10.1007/BF02552686
- Edwards BS, Edwards WD, Connolly DC, Edwards JE. Arterial-esophageal fistulae developing in patients with anomalies of the aortic arch system. *Chest*. 1984;86(5):732-735. doi: 10.1378/chest.86.5.732
- Gossot D, Nussaume O, Kitzis M, Cohen G, Chalaux G, Andreassian B. Fatal hematemesis due to erosion of a retro-esophageal right subclavian artery by an esophagogastric tube. Article in French. *Presse Med*. 1985;14(31):1655-1656.
- Kullnig P. Aneurysm of an aberrant right subclavian artery with bleeding into the esophageal wall. *Semin Roentgenol*. 1989;24(2):75-76. doi: 10.1016/0037-198x(89)90027-8
- Stone WM, Brewster DC, Moncure AC, Franklin DP, Cambria RP, Abbott WM. Aberrant right subclavian artery: varied presentations and management options. *J Vasc Surg*. 1990;11(6):812-817.
- Ikeda T, Yokota Y, Ando F, et al. A case of an aberrant subclavian artery-esophageal fistula due to prolonged nasogastric intubation. Article in Japanese. *Kyobu Geka*. 1991;44(12):1045-1047.
- Hirakata R, Hasuo K, Yasumori K, Yoshida K, Masuda K. Arterioenteric fistulae: diagnosis and treatment by angiography. *Clin Radiol*. 1991;43(5):328-330. doi: 10.1016/s0009-9260(05)80541-0
- Miller RG, Robie DK, Davis SL, et al. Survival after aberrant right subclavian artery-esophageal fistula: case report and literature review. *J Vasc Surg*. 1996;24(2):271-275. doi: 10.1016/s0741-5214(96)70103-9
- Singha NK, Hale SJ, Kuhlman JE. Arterio-esophageal communication from a ruptured aberrant right subclavian artery aneurysm. CT diagnosis. *Clin Imaging*. 1998;22(2):117-121. doi: 10.1016/s0899-7071(97)00072-7
- Feugier P, Lemoine L, Beaudoin N, Chevalier JM. Aberrant right subclavian arterioesophageal fistula: endovascular occlusion via a transbrachial approach. *Eur J Vasc Endovasc Surg*. 2002;23(1):77-78. doi: 10.1053/ejvs.2001.1512
- Lehmann B, Clemetson I, Fantin AC, et al. Arterioesophageal fistula secondary to rupture of an aberrant right subclavian artery aneurysm: a rare differential diagnosis in upper gastrointestinal bleeding. *Endoscopy*. 2006;38(7):762. doi: 10.1055/s-2006-925172
- Millar A, Rostom A, Rasuli P, Saloojee N. Upper gastrointestinal bleeding secondary to an aberrant right subclavian artery-esophageal fistula: a case report and review of the literature. *Can J Gastroenterol*. 2007;21(6):389-392. doi: 10.1155/2007/398213
- Magagna P, Abbiate N, Mansi G, et al. Endovascular treatment of aberrant right subclavian (lusorian) artery to oesophagus fistula: a case report. *Vasc Endovascular Surg*. 2008;42(4):394-396. doi: 10.1177/1538574408315993
- Fuentes S, Cano I, López M, et al. Arterial-esophageal fistula: a severe complication in children with cardiovascular abnormalities. *Pediatr Surg Int*. 2010;26(3):335-337. doi: 10.1007/s00383-009-2532-6
- Chapman JR, Sedghi S, Christie BD, Nakayama DK, Wynne JL. Aberrant right subclavian artery-esophageal fistula. *Am Surg*. 2010;76(12):1430-1432.

30. Situma M, Kubiak R, Numanoglu A, Wood R, Brooks A, Millar AJ. Near-fatal bleeding from an aberrant subclavian artery following colonic interposition for oesophageal atresia. *Pediatr Surg Int*. 2011;27(10):1131-1133. doi: 10.1007/s00383-011-2881-9
31. Jain KK, Braze AJ, Shapiro MA, Perez-Tamayo RA. Aberrant right subclavian artery-esophageal fistula and severe gastrointestinal bleeding after surgical correction of scimitar syndrome. *Tex Heart Inst J*. 2012;39(4):571-574.
32. Pop D, Venissac N, Nadeemy AS, Schneck AS, Aze O, Mouroux J. Lesson to be learned: beware of lusoria artery during transhiatal esophagectomy. *Ann Thorac Surg*. 2012;94(3):1010-1011. doi: 10.1016/j.athoracsur.2012.01.092
33. Takahashi S, Okada K, Orihashi K, Sueda T. Arterio-oesophageal fistula caused by aberrant right subclavian artery aneurysm. *Interact Cardiovasc Thorac Surg*. 2013;16(6):920-922. doi: 10.1093/icvts/ivt083
34. Lo A, Baird R, De Angelis P, et al. Arterioesophageal fistula after stenting for esophageal atresia. *J Pediatr Gastroenterol Nutr*. 2013;56(5):e30-e31. doi: 10.1097/MPG.0b013e31824ffd7f
35. Morisaki A, Hirai H, Sasaki Y, Hige K, Bito Y, Suehiro S. Aortoesophageal fistula after endovascular repair for aberrant right subclavian artery aneurysm. *Ann Thorac Cardiovasc Surg*. 2014;20 Suppl:790-793. doi: 10.5761/atcs.cr.12.02153
36. Hosn MA, Haddad F, El-Merhi F, Safadi B, Hallal A. Repair of an aberrant subclavian arterioesophageal fistula following esophageal stent placement. *World J Gastrointest Surg*. 2014;6(6):117-121. doi: 10.4240/wjgs.v6.i6.117
37. Joynt MR, Grifka RG. Spontaneous aberrant right subclavian arterio-oesophageal fistula in a previously healthy child. *Cardiol Young*. 2015;25(7):1425-1427. doi: 10.1017/S1047951114002388
38. Oliveira E, Anastácio M, Marques A. Aberrant right subclavian artery-esophageal fistula: massive upper gastrointestinal hemorrhage secondary to prolonged intubation. *Braz J Anesthesiol*. 2016;66(3):318-320. doi: 10.1016/j.bjane.2013.07.019
39. Watanabe M, Suzuki K, Fujinaga K, et al. Postmortem diagnosis of massive gastrointestinal bleeding in a patient with aberrant right subclavian artery-esophageal fistula. *Acute Med Surg*. 2015;3(2):139-142. doi: 10.1002/ams2.136
40. Kudose S, Pineda J, Saito JM, Dehner LP. Aberrant right subclavian artery-esophageal fistula in 20-year-old with VATER association. *J Pediatr Intensive Care*. 2017;6(2):127-131. doi: 10.1055/s-0036-1584812
41. Shires CB, Rohrer MJ. Anomalous right subclavian artery-esophageal fistulae. *Case Rep Vasc Med*. 2018;2018:7541904. doi: 10.1155/2018/7541904
42. Zheng S, Wang X, Chen S, et al. A pseudoaneurysm of aberrant right subclavian artery caused by esophageal stent placement because of esophageal stricture after endoscopic submucosal dissection. *Surg Laparosc Endosc Percutan Tech*. 2019;29(5):e69-e71. doi: 10.1097/SLE.0000000000000696
43. Merlo A, Farber M, Ohana E, Pascarella L, Crowner J, Long J. Aberrant right subclavian artery to esophageal fistula: a rare case and its management. *Ann Thorac Surg*. 2020;110(2):e85-e86. doi: 10.1016/j.athoracsur.2019.12.045
44. Jonker FHW, Schlosser FJV, Geirsson A, Sumpio BE, Moll FL, Muhs BE. Endograft collapse after thoracic endovascular aortic repair. *J Endovasc Ther*. 2010;17(6):725-734. doi: 10.1583/10-3130.1

This article meets the Accreditation Council for Graduate Medical Education and the American Board of Medical Specialties Maintenance of Certification competencies for Patient Care and Medical Knowledge.

©2021 by the author(s); licensee Ochsner Journal, Ochsner Clinic Foundation, New Orleans, LA. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (creativecommons.org/licenses/by/4.0/legalcode) that permits unrestricted use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.