

Cross-Sectional Survey of Smoking Patterns During the COVID-19 Pandemic in a Tobacco Cessation and Lung Cancer Screening Program

Maria C. Mejia, MD, MPH, FACPM,¹ Roger Zoorob, MD, MPH,¹ Robert S. Levine, MD,^{1,2} Xiaofan Huang, MS,³ Charles H. Hennekens, MD, DrPH²

¹Department of Family and Community Medicine, Baylor College of Medicine, Houston, TX ²Charles E. Schmidt College of Medicine, Florida Atlantic University, Boca Raton, FL ³Dan L. Duncan Institute for Clinical and Translational Research, Baylor College of Medicine, Houston, TX

Background: Coronavirus disease 2019 (COVID-19) produces a wide array of deleterious consequences, some of which are unintended. Data are sparse on whether, and if so, how, current cigarette smoking habits are affected by COVID-19. We describe changes to smoking habits and their correlates during the COVID-19 pandemic among participants in a tobacco cessation and lung cancer screening program.

Methods: Between June and October 2020, we conducted a cross-sectional survey of a convenience sample of 150 participants in a lung cancer screening and tobacco cessation program. The survey consisted of 3 parts: (1) changes in tobacco use, (2) impact and coping strategies toward COVID-19, and (3) COVID-19 exposure and use of protective measures. Demographic variables included age, sex, race/ethnicity, and marital status.

Results: All 150 participants who were contacted agreed to participate in this cross-sectional survey. The statistically significant correlates of increased tobacco use were high uncertainty about the future ($P < 0.001$), loneliness because of social distancing or self-isolating ($P < 0.001$), anger or frustration with how the pandemic has disrupted daily life ($P < 0.001$), boredom resulting from inability to work or engage in regular daily activities/routines ($P < 0.001$), desire to cope using alcohol or drugs ($P = 0.002$), sadness or feelings of hopelessness ($P = 0.003$), and worry or fear about challenges to securing basic needs such as groceries or medication ($P < 0.001$). In contrast, those who smoked less were more likely to practice social distancing ($P = 0.002$) and use protective measures ($P = 0.005$).

Conclusion: Among those who decreased or stopped smoking, correlates included greater use of protective measures for COVID-19, including social distancing and testing. These data may aid healthcare providers to identify and provide counsel to cigarette smokers at greater risks for increasing tobacco consumption during stresses such as COVID-19.

Keywords: COVID-19, smoking, smoking cessation, tobacco cessation, tobacco smoking

Address correspondence to Maria C. Mejia, MD, MPH, FACPM, Department of Family and Community Medicine, Baylor College of Medicine, 3701 Kirby Dr., Ste. 600, Houston, TX 77098. Tel: (713) 798-4735. Email: maria.mejia@bcm.edu

INTRODUCTION

Since 1965, the prevalence of cigarette smoking has markedly decreased in the United States.¹ Despite the remarkable decline, cigarette smoking causes hundreds of thousands of potentially avoidable premature deaths in the United States each year, principally from lung cancer, cardiovascular disease, and chronic respiratory disease.¹ While coronavirus disease 2019 (COVID-19) has caused more than 944,000 premature deaths since January 2020, the US Centers for Disease Control and Prevention estimates that tobacco-related causes account for more than 480,000 deaths per year in the United States.^{1,2}

Tobacco consumption among US smokers has increased by more than 30% during the COVID-19 pandemic.³ COVID-19 has also adversely impacted screening,⁴ diagnosis,⁵ and survival from lung cancer.⁶ Moreover, concerns have been raised about adverse effects of the COVID-19 pandemic on smokers trying to quit in the United States.⁷

To the best of our knowledge, however, data are sparse about the possible interrelationships and characteristics of cigarette smokers who have increased and decreased their habits during the COVID-19 pandemic. In this report, we describe changes to smoking habits and correlates of increases and decreases during the COVID-19 pandemic, including key indicators of psychological distress among

participants enrolled in a tobacco cessation and lung cancer screening program.

METHODS

Design

Between June and October 2020, we conducted a cross-sectional survey of a convenience sample of 150 participants in a lung cancer screening and tobacco cessation program. Participants were eligible for the program if they were 55 to 80 years of age, had a tobacco history of at least 30 pack-years, and were currently smoking or had quit smoking within the prior 15 years.⁸ The program involved 6 to 8 visits, including an initial face-to-face contact lasting 45 to 60 minutes and follow-up in person or by phone. As the US COVID-19 epidemic increased markedly, all visits transitioned to phone contacts, and educational materials were mailed to participants along with nicotine replacement therapies if indicated. Each participant was asked about participating in the survey during regular phone follow-up until 150 participants were enrolled. Data were collected by 3 trained interviewers whose backgrounds included fact-checking and verification.

Our sample size of 150 provided sufficient statistical power to detect small to moderate differences in smoking habits.⁹ Findings are reported in accordance with the Strengthening of the Reporting of Observational Studies in Epidemiology (STROBE) guidelines.¹⁰ The study was approved by the Institutional Review Board of the Baylor College of Medicine.

Measures

The survey consisted of 3 parts: (1) changes in tobacco use, (2) impact and coping strategies toward COVID-19, and (3) COVID-19 exposure and use of protective measures. Demographic variables included age, sex, race/ethnicity, and marital status. The survey (Appendix) included 28 questions adapted from the CoRonavluS Health Impact Survey¹¹ and the US Centers for Disease and Control Prevention guidance for COVID-19.¹² Tobacco use change during the COVID-19 pandemic was classified as increased, unchanged, decreased, varied day to day with no clear pattern, and not applicable/former smoker. Coping strategies and mental health questions were answered on a 5-point Likert scale from least (1) to greatest (5). We included 4 questions about the clinical symptoms of COVID-19 and 3 questions about protective strategies. Five questions referred to mitigation practices, also scored on a 5-point Likert scale.

Data Analysis

We first explored whether significant differences were found among those who increased, decreased, or continued their same smoking patterns. Categorical variables are summarized by frequencies and percentages. Continuous variables are summarized by medians with interquartile ranges. We stratified the data by changes in smoking use overall and by race/ethnicity and tested for statistical significance using Kruskal-Wallis or chi-squared tests. To explore interrelationships of changes in smoking with categorical baseline characteristics, we used paired chi-squared tests with Holm *P* value adjustments to explore which smoking use change and race/ethnicity group were significantly different from other categories. To explore continuous variables, we

Table 1. Participant Demographics, n=150

Variable	Value
Age, years, mean ± SD	60.7 ± 4.5
Sex	
Female	53 (35)
Male	97 (65)
Race	
Black	73 (49)
White	64 (43)
Asian/Pacific Islander	13 (9)
Ethnicity	
Non-Hispanic	117 (78)
Hispanic	33 (22)
Marital status ^a	
Single	72 (51)
Married	41 (29)
Divorced	24 (17)
Widowed	5 (4)

Note: Data are presented as n (%) unless otherwise indicated.

^an=142.

used pairwise Wilcoxon rank sum tests with Holm *P* value adjustments. We considered 2-sided *P* values at <0.05 as significant.

RESULTS

Among the 150 participants, 50 (33%) reported changes in tobacco use, 60 (40%) reported no changes, and 40 (27%) were former smokers. No former smokers reported relapsing. The mean age was 60.7 ± 4.5 years, and 97 participants (65%) were male (Table 1). Participants who reported increases in smoking had high levels of uncertainty about the future ($P<0.001$) (Figure 1A), loneliness as a result of social distancing or self-isolating ($P<0.001$) (Figure 1B), anger or frustration with how the pandemic has disrupted daily life ($P<0.001$) (Figure 2A), boredom because of being unable to work or engage in regular daily activities/routines ($P=0.001$) (Figure 2B), a desire to cope using alcohol or drugs ($P=0.002$) (Figure 3A), sadness or feelings of hopelessness ($P=0.003$) (Figure 3B), and worry or fear about challenges to securing basic needs such as groceries or medication ($P<0.001$) (Figure 4A). In contrast, those who smoked less were more likely to practice social distancing ($P=0.002$) (Figure 4B) and practice other protective measures ($P=0.005$) (Figure 5A).

Table 2 shows survey responses stratified by tobacco usage. The numbers of participants tested for COVID-19 were significantly different between the groups ($P=0.022$). We found no statistically significant difference between tobacco use status and current housing, whether the participant had perceived control over contracting COVID-19, or source of information about COVID-19.

Table 3 shows survey responses stratified by race/ethnicity. Among the 150 participants, 48.7% were Black, 20.7% were non-Hispanic White, 22% were Hispanic, and 8.7% were Asian/Pacific Islander. We found significant

Table 2. Survey Questions Stratified by Tobacco Usage, n=110

Survey Question	Change in Tobacco Usage			P Value
	Increased, n=31	Stayed the Same, n=60	Decreased, n=19	
Tobacco usage has changed during the COVID-19 pandemic, n (%)	31 (100.0)	0 (0.0)	19 (100.0)	<0.001
COVID-19 pandemic has affected your desire to participate in our LDCT lung cancer screening program, n (%)	4 (12.9)	3 (5.0)	2 (10.5)	0.393
Coping strategies and mental health, median [IQR]				
Uncertainty about the future	4.00 [4.00, 5.00]	3.00 [1.00, 4.00]	2.00 [1.00, 3.00]	<0.001
Loneliness as a result of social distancing/self-isolating	4.00 [3.00, 4.50]	2.00 [1.00, 3.00]	1.00 [1.00, 2.00]	<0.001
Anger or frustration with how the pandemic has disrupted your daily life	4.00 [3.50, 4.00]	3.00 [2.00, 4.00]	1.00 [1.00, 5.00]	<0.001
Boredom due to being unable to work or engage in regular daily activities/routines	4.00 [3.00, 5.00]	3.00 [2.00, 4.00]	3.00 [1.00, 3.00]	0.001
Desire to cope using alcohol or drugs	2.00 [1.00, 3.00]	2.00 [1.00, 2.00]	1.00 [1.00, 1.00]	0.002
Sadness or feelings of hopelessness	3.00 [2.50, 4.00]	2.00 [1.00, 3.00]	1.00 [1.00, 2.75]	0.003
Worry or fear about challenges to securing basic needs such as groceries or medication	4.00 [3.00, 4.50]	3.00 [1.00, 4.00]	3.00 [1.00, 4.00]	<0.001
Health management, n (%)				
Have a serious or chronic health condition that requires medication and management at home	12 (38.7)	20 (33.3)	12 (63.2)	0.068
Effects on housing and employment, n (%)				
I live alone	7 (22.6)	18 (30.0)	5 (26.3)	0.749
I live with spouse/significant other	11 (35.5)	21 (35.0)	9 (47.4)	0.634
I live with another family member	7 (22.6)	16 (26.7)	4 (21.1)	0.846
I live with a non-family member	6 (19.4)	5 (8.3)	1 (5.3)	0.191
COVID-19 affected my housing situation	6 (19.4)	5 (8.3)	1 (5.3)	0.191
COVID-19 affected my employment status	9 (29.0)	15 (25.0)	5 (26.3)	0.911
Exposure and preventive measures				
COVID-19 symptoms in the past 2 weeks, n (%)				
Fever	0 (0.0)	1 (1.7)	0 (0.0)	0.657
Dry cough	0 (0.0)	2 (3.3)	0 (0.0)	0.428
Runny nose	0 (0.0)	1 (1.7)	0 (0.0)	0.657
Body aches	0 (0.0)	1 (1.7)	1 (5.3)	0.398
Upset stomach	0 (0.0)	0 (0.0)	0 (0.0)	NA
Loss of smell/taste	0 (0.0)	0 (0.0)	0 (0.0)	NA
Fatigue, tiredness	1 (3.2)	1 (1.7)	1 (5.3)	0.689
None of the above	30 (96.8)	56 (93.3)	18 (94.7)	0.790
Had close contact with an individual with flu-like symptoms or a confirmed diagnosis of COVID-19 in the past 3 weeks, n (%)				
Yes	2 (6.5)	1 (1.7)	0 (0.0)	
No	27 (87.1)	55 (91.7)	16 (84.2)	
Unsure	2 (6.5)	4 (6.7)	3 (15.8)	
Have been tested for COVID-19, n (%)	4 (12.9)	14 (23.3)	9 (47.4)	0.022
Believed might have COVID-19 but did not get tested, n (%)	1 (3.2)	5 (8.3)	0 (0.0)	0.662

Table 2. Continued

Survey Question	Change in Tobacco Usage			P Value
	Increased, n=31	Stayed the Same, n=60	Decreased, n=19	
Reside in area under a “shelter-in-place” or “stay-at-home” order, n (%)	31 (100.0)	59 (98.3)	18 (94.7)	–
Avoided crowded places whenever possible, n (%)	30 (96.8)	58 (96.7)	19 (100.0)	0.724
Wore a mask when leaving the house, n (%)	30 (96.8)	60 (100.0)	19 (100.0)	0.276
Worried about you or individuals in your household getting COVID-19, median [IQR]	4.00 [2.00, 4.00]	2.50 [2.00, 4.00]	2.00 [1.00, 5.00]	0.081
Practice social distancing, median [IQR]	4.00 [4.00, 5.00]	4.00 [4.00, 5.00]	5.00 [5.00, 5.00]	0.002
Practice protective measures such as hand washing, disinfecting household surfaces, using hand sanitizer, median [IQR]	4.00 [3.50, 5.00]	4.00 [4.00, 5.00]	5.00 [5.00, 5.00]	0.005
Control you feel like you have over whether or not you or household might contract COVID-19, median [IQR]	3.00 [3.00, 4.00]	4.00 [3.00, 5.00]	4.00 [3.00, 5.00]	0.094
Feel equipped with enough knowledge to protect yourself and your household from COVID-19, median [IQR]	3.00 [3.00, 4.00]	4.00 [3.00, 5.00]	4.5 [3.25, 5.00]	0.064
Source of COVID-19 information and updates, n (%)				
National news outlet (FOX, CNN, MSNBC, etc.)	20 (64.5)	37 (61.7)	13 (68.4)	0.861
Local news on television	27 (87.1)	53 (88.3)	17 (89.5)	0.967
Newspaper or local print media	1 (3.2)	3 (5.0)	2 (10.5)	0.530
Directly from family or friends	8 (25.8)	14 (23.3)	9 (47.4)	0.120
A healthcare provider	2 (6.5)	6 (10.0)	4 (21.1)	0.260
Social media (Facebook, Instagram, Twitter)	3 (9.7)	12 (20.0)	5 (26.3)	0.289
Messaging platform (WhatsApp, Facebook)	0 (0.0)	1 (1.7)	2 (10.5)	0.065
Other	2 (6.5)	7 (11.7)	1 (5.3)	0.583

Note: Responses reported as median [interquartile range (IQR)] were scored according to a Likert scale ranging from 1 (least) to 5 (greatest). COVID-19, coronavirus disease 2019; LDCT, low dose computed tomography.

differences by race/ethnicity in the effect of COVID-19 on employment status ($P=0.042$). The levels of “How worried are you about you or individuals in your household getting COVID-19?” differed significantly by race/ethnicity ($P=0.004$), and Figure 5B shows that Asian participants were more worried than the participants in other racial/ethnic groups. The number of Hispanic participants who avoided crowded places was significantly lower than the other groups ($P=0.029$). Compared to other groups, a significantly lower number of Hispanic participants received COVID-19 information from a national news outlet ($P=0.033$), and more Hispanics received information directly from family or friends ($P=0.003$). We found no statistically significant differences by race/ethnicity with respect to change in tobacco usage, desire to participate in the screening/tobacco control program, uncertainty about the future, loneliness, anger or frustration about how the pandemic has disrupted daily life, boredom, desire to cope with alcohol or drugs, feelings of sadness or hopelessness, or worry or fear about challenges to securing basic needs.

DISCUSSION

Among 150 participants in a tobacco cessation and lung cancer screening program during the COVID-19 pandemic,

we found statistically significant and potentially clinically important differences between those who increased and decreased tobacco use. Among current smokers, 28.2% (31/110) reported increased tobacco use, 17.3% (19/110) decreased, and 54.5% (60/110) no change. In addition, there were no reports of relapse among former smokers.

We found correlates of increased tobacco use related to coping strategies and mental health such as high uncertainty about the future, loneliness as a result of social distancing, anger or frustration with how the pandemic has disrupted daily life, boredom because of being unable to work or engage in regular daily activities/routines, desire to cope using alcohol or drugs, sadness or feelings of hopelessness, and worry or fear about challenges to securing basic needs such as groceries or medication. In contrast, those who smoked less were more likely to practice social distancing and other protective measures.

The current data are compatible with previous reports of a bidirectional effect of COVID-19 observed in smokers: a reduction in or cessation of smoking in some but an increase in others.^{13,14} In an online survey of 1,491 adults in Australia, all aspects of psychological distress—specifically depression, anxiety, and stress—were significantly correlated with health behavior.¹⁵ Among 172 smokers completing the

Table 3. Survey Questions Stratified by Race/Ethnicity, n=150

Survey Question	Race/Ethnicity				P Value
	Non-Hispanic White, n=31	Hispanic, n=33	Black, n=73	Asian/Pacific Islander, n=13	
Tobacco usage has changed during the pandemic, n (%)	12 (38.7)	9 (27.3)	25 (34.2)	4 (30.8)	0.600
Way your tobacco usage changed during the COVID-19 pandemic, n (%)	n=23	n=20	n=58	n=9	0.748
Increased	8 (34.8)	5 (25.0)	17 (29.3)	1 (11.1)	
Decreased	4 (17.4)	4 (20.0)	8 (13.8)	3 (33.3)	
Stayed the same	11 (47.8)	11 (55.0)	33 (56.9)	5 (55.6)	
COVID-19 pandemic has affected your desire to participate in our LDCT lung cancer screening program, n (%)	2 (6.5)	2 (6.1)	8 (11.0)	0 (0.0)	0.519
Coping strategies and mental health, median [IQR]					
Uncertainty about the future	3.00 [3.00, 4.00]	4.00 [2.00, 5.00]	3.00 [1.00, 4.00]	4.00 [2.00, 4.00]	0.467
Loneliness as a result of social distancing/self-isolating	3.00 [2.00, 4.00]	3.00 [2.00, 3.00]	2.00 [1.00, 4.00]	2.00 [1.00, 3.00]	0.484
Anger or frustration with how the pandemic has disrupted your daily life	3.00 [2.00, 4.00]	4.00 [2.00, 4.00]	3.00 [2.00, 4.00]	3.00 [2.00, 3.00]	0.489
Boredom due to being unable to work or engage in regular daily activities/routines	3.00 [2.00, 4.00]	3.00 [2.00, 4.00]	3.00 [2.00, 4.00]	3.00 [2.00, 3.00]	0.855
Desire to cope using alcohol or drugs	1.00 [1.00, 2.00]	1.00 [1.00, 2.00]	1.00 [1.00, 2.00]	1.00 [1.00, 1.00]	0.172
Sadness or feelings of hopelessness	2.00 [2.00, 3.00]	2.00 [1.00, 3.00]	2.00 [1.00, 3.00]	2.00 [1.75, 4.25]	0.839
Worry or fear about challenges to securing basic needs such as groceries or medication	3.00 [2.00, 4.00]	4.00 [1.00, 4.00]	3.00 [2.00, 4.00]	3.00 [3.00, 4.00]	0.901
Health management, n (%)					
Have a serious or chronic health condition that requires medication and management at home	13 (41.9)	9 (27.3)	30 (41.1)	3 (23.1)	0.351
Effects on housing and employment, n (%)					
I live alone	10 (32.3)	4 (12.1)	31 (42.5)	0 (0.0)	0.001
I live with spouse/significant other	11 (35.5)	20 (60.6)	19 (26.0)	11 (84.6)	<0.001
I live with another family member	6 (19.4)	7 (21.2)	17 (23.3)	2 (15.4)	0.916
I live with a non-family member	4 (12.9)	2 (6.1)	6 (8.2)	0 (0.0)	0.509
COVID-19 affected my housing situation	4 (12.9)	2 (6.1)	9 (12.5)	1 (7.7)	0.738
COVID-19 affected my employment status	5 (16.1)	10 (30.3)	15 (20.5)	7 (53.8)	0.042
Exposure and preventive measures					
COVID-19 symptoms in the past 2 weeks, n (%)					
Fever	0 (0.0)	0 (0.0)	1 (1.4)	0 (0.0)	0.786
Dry cough	0 (0.0)	0 (0.0)	2 (2.7)	0 (0.0)	0.544
Runny nose	0 (0.0)	0 (0.0)	1 (1.4)	0 (0.0)	0.786
Body aches	0 (0.0)	0 (0.0)	2 (2.7)	0 (0.0)	0.544
Upset stomach	0 (0.0)	0 (0.0)	1 (1.4)	0 (0.0)	0.786
Loss of smell/taste	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	N/A
Fatigue, tiredness	0 (0.0)	0 (0.0)	3 (4.1)	0 (0.0)	0.358
None of the above	31 (100.0)	33 (100.0)	66 (90.4)	13 (100.0)	0.052

Table 3. Continued

Survey Question	Race/Ethnicity				P Value
	Non-Hispanic White, n=31	Hispanic, n=33	Black, n=73	Asian/Pacific Islander, n=13	
Had close contact with an individual with flu-like symptoms or a confirmed diagnosis of COVID-19 in the past 3 weeks, n (%)					0.816
Yes	1 (3.2)	1 (3.0)	3 (4.1)	0 (0.0)	
No	26 (83.9)	31 (93.9)	62 (84.9)	12 (92.3)	
Unsure	4 (12.9)	1 (3.0)	8 (11.0)	1 (7.7)	
Have been tested for COVID-19, n (%)	7 (22.6)	4 (12.1)	23 (31.5)	2 (15.4)	0.147
Believed might have had COVID-19 but did not get tested, n (%)	2 (6.5)	2 (6.1)	3 (4.1)	0 (0.0)	0.786
Reside in an area under a "shelter-in-place" or "stay-at-home" order, n (%)	30 (96.8)	33 (100.0)	70 (95.9)	13 (100.0)	–
Avoided crowded places whenever possible, n (%)	30 (96.8)	28 (84.8)	72 (98.6)	12 (92.3)	0.029
Wore a mask when leaving the house, n (%)	31 (100.0)	32 (97.0)	73 (100.0)	13 (100.0)	0.312
Worried about you or individuals in your household getting COVID-19, median [IQR]	3.00 [2.00, 4.00]	3.00 [2.00, 4.00]	2.00 [2.00, 4.00]	4.00 [3.00, 5.00]	0.004
Practice social distancing, median [IQR]	5.00 [4.00, 5.00]	4.00 [4.00, 5.00]	5.00 [4.00, 5.00]	4.00 [4.00, 5.00]	0.602
Practice protective measures such as hand washing, disinfecting household surfaces, using hand sanitizer, median [IQR]	5.00 [4.00, 5.00]	4.00 [4.00, 5.00]	5.00 [4.00, 5.00]	4.00 [4.00, 5.00]	0.677
Control you feel you have over whether or not you or household might contract COVID-19, median [IQR]	4.00 [3.00, 4.00]	3.00 [2.00, 4.00]	4.00 [3.00, 4.00]	3.00 [3.00, 4.00]	0.394
Feel equipped with enough knowledge to protect yourself and your household from COVID-19, median [IQR]	4.00 [3.00, 4.00]	3.00 [3.00, 4.00]	4.00 [3.00, 5.00]	3.50 [3.00, 5.00]	0.068
Source of COVID-19 information and updates, n (%)					
National news outlet (FOX, CNN, MSNBC, etc.)	22 (71.0)	15 (45.5)	54 (74.0)	9 (69.2)	0.033
Local news on television	30 (96.8)	31 (93.9)	62 (84.9)	11 (84.6)	0.229
Newspaper or local print media	3 (9.7)	2 (6.1)	2 (2.7)	0 (0.0)	0.371
Directly from family or friends	5 (16.1)	16 (48.5)	13 (17.8)	5 (38.5)	0.003
A healthcare provider	2 (6.5)	4 (12.1)	9 (12.3)	1 (7.7)	0.805
Social media (Facebook, Instagram, Twitter)	7 (22.6)	4 (12.1)	9 (12.3)	2 (15.4)	0.563
Messaging platform (WhatsApp, Facebook)	0 (0.0)	2 (6.1)	2 (2.7)	0 (0.0)	0.445
Other	3 (9.7)	1 (3.0)	6 (8.2)	3 (23.1)	0.188

Note: Responses reported as median [interquartile range (IQR)] were scored according to a Likert scale ranging from 1 (least) to 5 (greatest). COVID-19, coronavirus disease 2019; LDCT, low dose computed tomography.

survey, 49.9% reported an increase in smoking behavior, while 16.3% reported a reduction.¹⁵ An online survey of 4,005 French adults during COVID-19 included 1,062 regular smokers; 231 (21.8%) reported that they increased their intake, while 177 (16.7%) reportedly decreased intake.¹⁶ The study authors suggested that the threat of contracting COVID-19 may have motivated some smokers to improve, while boredom and restrictions in movement may have had the opposite effect in others. In a survey from Italy, smokers reported increased tobacco use, increased food intake, and

changes (positive and negative) in sleep quality during home confinement.¹⁷ High stress levels and the need to adapt to prolonged stays at home, particularly among fully employed individuals, have had significant correlations during COVID-19.¹⁸

Smoking cessation programs struggle to achieve results. A report by Lang and Yakhkind showed that while the pandemic provides an opportunity to adapt and expand smoking cessation services, potential challenges involve the increased need for behaviorists because of widespread




Figure 1. (A) Uncertainty about the future. Participants with increased tobacco use had more uncertainty about the future compared to those with unchanged (median 4 vs 3, $P<0.001$) or decreased tobacco use (median 4 vs 2, $P=0.002$). **(B) Loneliness as a result of social distancing.** Participants with increased tobacco use were more likely to report loneliness as a result of social distancing compared to those with unchanged (median 4 vs 2, $P<0.001$) or decreased tobacco use (median 4 vs 1, $P<0.001$). NA, not applicable.

stress brought on, in part, by isolation from friends and family.¹⁹ Some smokers may continue to smoke because of their erroneous belief that tobacco is a protective factor against COVID-19.¹⁹ Such erroneous beliefs may

lead to decreases in enrollment in smoking cessation programs.¹⁴

In a survey of smokers in Australia and the United Kingdom, 45% of respondents wanted more information about




Figure 2. (A) Anger or frustration with how the pandemic has disrupted your daily life. Participants with increased tobacco use reported more anger or frustration with how the pandemic has disrupted their daily life compared to those with unchanged tobacco use (median 4 vs 3, $P<0.001$). **(B) Boredom due to being unable to work or engage in regular daily activities/routines.** Participants with increased tobacco use reported more boredom compared to those with unchanged (median 4 vs 3, $P=0.001$) or decreased tobacco use (median 4 vs 3, $P=0.001$). NA, not applicable.




Figure 3. (A) Desire to cope using alcohol or drugs. Participants with decreased tobacco use had less desire to cope using alcohol or drugs compared to those with unchanged (median 1 vs 2, $P=0.008$) or increased tobacco use (median 1 vs 2, $P=0.002$). (B) Sadness, or feelings of hopelessness. Participants with increased tobacco use had more sadness or feelings of hopelessness compared to those with unchanged (median 3 vs 2, $P=0.006$) or decreased tobacco use (median 3 vs 1, $P=0.003$). NA, not applicable.

smoking and COVID-19 risk.²⁰ The most popular sources of information identified by respondents were government departments (59%) and physicians (47%), with the preferred delivery platforms being television (61%), online news sources (36%), social media (31%), and e-mail (31%).²⁰ This finding contrasts with the data from our study that indicates local and national news outlets were the most common

sources of information about the pandemic for respondents, with Hispanics relying more on friends and family members for information than Asians/Pacific Islanders, Blacks, and non-Hispanic Whites.

This study has several limitations that merit consideration. First are the restrictions on generalizability. Participants were tobacco users recruited from a lung cancer screening




Figure 4. (A) Worry/fear about challenges to securing basic needs such as groceries or medication. Participants with decreased tobacco use had more worry or fear about challenges to securing basic needs compared to those with unchanged tobacco use (median 4 vs 3, $P<0.001$). (B) How much social distancing do you practice? Participants with decreased tobacco use practiced more social distancing than those with unchanged (median 5 vs 4, $P=0.012$) or increased tobacco use (median 5 vs 4, $P=0.002$). NA, not applicable.




Figure 5. (A) How often do you practice protective measures such as hand washing, disinfecting household surfaces, using hand sanitizer? Participants with decreased tobacco use practiced protective measures more often than those with unchanged (median 5 vs 4, $P=0.027$) or increased tobacco use (median 5 vs 4, $P=0.005$). (B) How worried are you about you or individuals in your household getting COVID-19? Participants who were Asian worried more about themselves or individuals in their household getting COVID-19 compared to patients who were non-Hispanic White (median 4 vs 3, $P=0.025$), Hispanic (median 4 vs 3, $P=0.037$), or Black (median 4 vs 2, $P=0.002$). NA, not applicable.

program who had volunteered to participate in a tobacco cessation program. In addition, all responses were self-reported and may be subject to recall bias. Also, as with any cross-sectional survey, the data are useful to formulate but not test hypotheses.^{21,22}

One unique strength of this cross-sectional survey is that all 150 volunteers approached agreed to participate. Another strength is that many hypotheses can be formulated from these descriptive data regarding possible behavioral and other factors that led some individuals to increase and others to decrease their smoking habits during COVID-19. Tests of such hypotheses will require analytic epidemiologic research designed a priori to do so.

CONCLUSION

These descriptive data show statistically significant differences between those who increased and decreased tobacco use during COVID-19 among volunteers who enrolled in a tobacco cessation and lung cancer screening program in Houston, Texas. Among those who decreased or stopped smoking, correlates included greater use of protective measures for COVID-19 such as social distancing and testing. These data may aid healthcare providers to identify and provide counsel to cigarette smokers at greater risk for increasing tobacco consumption during stresses such as the COVID-19 pandemic.

ACKNOWLEDGMENTS

This work was supported by the Cancer Prevention and Research Institute of Texas, grant number PP180016. Dr Mejia, Professors Zoorob and Levine, and Ms Huang have no disclosures. Professor Hennekens reports that he serves as

an independent scientist in an advisory role to investigators as chair of data monitoring committees for Amgen, British Heart Foundation, Cadila, Canadian Institutes of Health Research, DalCor, and Regeneron, and to the Collaborative Institutional Training Initiative, legal counsel for Pfizer, the US Food and Drug Administration, and UpToDate. Professor Hennekens receives royalties for authorship or editorship of 3 textbooks and as co-inventor on patents for inflammatory markers of cardiovascular disease that are held by Brigham and Women’s Hospital; has an investment management relationship with the West-Bacon Group within SunTrust Investment Services, which has discretionary investment authority; and does not own any common or preferred stock in any pharmaceutical or medical device company.

The authors are grateful to the survey field and data management team members for their dedication and hard work: April Walker, MA, Adriana Rangel, MPH, and Kimberly Hinh, MPH. The authors are indebted to Kristen Staggers, MS, for her expert assistance with the data analysis.

REFERENCES

1. Creamer MR, Wang TW, Babb S, et al. Tobacco product use and cessation indicators among adults – United States, 2018. *MMWR Morb Mortal Wkly Rep.* 2019;68(45):1013-1019. doi: 10.15585/mmwr.mm6845a2
2. COVID Data Tracker. Centers for Disease Control and Prevention. Accessed February 28, 2022. covid.cdc.gov/covid-data-tracker
3. Caponnetto P, Inguscio L, Saitta C, Maglia M, Benfatto F, Polosa R. Smoking behavior and psychological dynamics during COVID-19 social distancing and stay-at-home policies: a survey. *Health Psychol Res.* 2020;8(1):9124. doi: 10.4081/hpr.2020.9124

4. Van Haren RM, Delman AM, Turner KM, et al. Impact of the COVID-19 pandemic on lung cancer screening program and subsequent lung cancer. *J Am Coll Surg.* 2021;232(4):600-605. doi: 10.1016/j.jamcollsurg.2020.12.002
5. Kaufman HW, Chen Z, Niles J, Fesko Y. Changes in the number of US patients with newly identified cancer before and during the coronavirus disease 2019 (COVID-19) pandemic. *JAMA Netw Open.* 2020;3(8):e2017267. doi: 10.1001/jamanetworkopen.2020.17267. Erratum in: *JAMA Netw Open.* 2020;3(9):e2020927. doi: 10.1001/jamanetworkopen.2020.20927
6. Chavez-MacGregor M, Lei X, Zhao H, Scheet P, Giordano SH. Evaluation of COVID-19 mortality and adverse outcomes in US patients with or without cancer. *JAMA Oncol.* 2022;8(1):69-78. doi: 10.1001/jamaoncol.2021.5148
7. Jaklevic MC. COVID-19 and the "lost year" for smokers trying to quit. *JAMA.* 2021;325(19):1929-1930. doi: 10.1001/jama.2021.5601
8. Blackmon SH, Feinglass SR. The United States Preventive Services Task Force recommendations for lung cancer screening. *Thorac Surg Clin.* 2015;25(2):199-203. doi: 10.1016/j.thorsurg.2014.12.004
9. Leeman J, Birken SA, Powell BJ, Rohweder C, Shea CM. Beyond "implementation strategies": classifying the full range of strategies used in implementation science and practice. *Implement Sci.* 2017;12(1):125. doi: 10.1186/s13012-017-0657-x
10. Vandembroucke JP, von Elm E, Altman DG, et al. Strengthening the Reporting of Observational Studies in Epidemiology (STROBE): explanation and elaboration. *Ann Intern Med.* 2007;147(8):W163-W194. doi: 10.7326/0003-4819-147-8-200710160-00010-w1
11. Merikangas K, Milham M, Stringaris A, Bromet E, Colcombe S, Zipunnikov V. The Coronavirus Health Impact Survey (CRISIS). Accessed February 18, 2022. www.crisissurvey.org/
12. COVID-19 Guidance Documents. Centers for Disease and Control Prevention. Accessed April 30, 2020. www.cdc.gov/coronavirus/2019-ncov/communication/guidance-list.html?Sort=Date%3A%3Adesc&Audience=General%20Public&Topic=Prevention%20%26%20Infection%20Control
13. Bommele J, Hopman P, Walters BH, et al. The double-edged relationship between COVID-19 stress and smoking: implications for smoking cessation. *Tob Induc Dis.* 2020;18:63. doi: 10.18332/tid/125580
14. Klemperer EM, West JC, Peasley-Miklus C, Villanti AC. Change in tobacco and electronic cigarette use and motivation to quit in response to COVID-19. *Nicotine Tob Res.* 2020;22(9):1662-1663. doi: 10.1093/ntr/ntaa072
15. Stanton R, To QG, Khalesi S, et al. Depression, anxiety and stress during COVID-19: associations with changes in physical activity, sleep, tobacco and alcohol use in Australian adults. *Int J Environ Res Public Health.* 2020;17(11):4065. doi: 10.3390/ijerph17114065
16. Constant A, Conserve DF, Gallopel-Morvan K, Raude J. Socio-cognitive factors associated with lifestyle changes in response to the COVID-19 epidemic in the general population: results from a cross-sectional study in France. *Front Psychol.* 2020;11:579460. doi: 10.3389/fpsyg.2020.579460
17. Canello R, Soranna D, Zambra G, Zambon A, Invitti C. Determinants of the lifestyle changes during COVID-19 pandemic in the residents of northern Italy. *Int J Environ Res Public Health.* 2020;17(17):6287. doi: 10.3390/ijerph17176287
18. Sidor A, Rzymiski P. Dietary choices and habits during COVID-19 lockdown: experience from Poland. *Nutrients.* 2020;12(6):1657. doi: 10.3390/nu12061657
19. Lang AE, Yakhkind A. Coronavirus disease 2019 and smoking: how and why we implemented a tobacco treatment campaign. *Chest.* 2020;158(4):1770-1776. doi: 10.1016/j.chest.2020.06.013
20. Pettigrew S, Jun M, Roberts I, Bullen C, Nallaiah K, Rodgers A. Preferences for tobacco cessation information and support during Covid-19. *J Addict Med.* 2020;14(6):e362-e365. doi: 10.1097/ADM.0000000000000743
21. Hennekens CH, DeMets D. Statistical association and causation: contributions of different types of evidence. *JAMA.* 2011;305(11):1134-1136. doi: 10.1001/jama.2011.322
22. Hennekens CH, Buring JE. *Epidemiology in Medicine.* Little, Brown and Company; 1987.

This article meets the Accreditation Council for Graduate Medical Education and the American Board of Medical Specialties Maintenance of Certification competencies for Patient Care, Medical Knowledge, and Practice-Based Learning and Improvement.

©2022 by the author(s); licensee Ochsner Journal, Ochsner Clinic Foundation, New Orleans, LA. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (creativecommons.org/licenses/by/4.0/legalcode) that permits unrestricted use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.



APPENDIX

Confidential

Page 1 of 3

COVID-19 Survey

Date _____

Participant's ID _____

Interviewer: _____

Tobacco Usage

Has your tobacco usage changed in any way (increased, decreased) during/after the COVID-10 pandemic? Yes No

In what way has your tobacco usage changed during/after the COVID-10 pandemic?

- My tobacco usage has increased
- My tobacco usage has stayed the same
- My tobacco usage has decreased
- My tobacco usage has varied from day to day with no clear pattern
- Not applicable/former smoker

Has the COVID-19 pandemic affected your desire to participate in our LDCT lung cancer screening program? Yes No

If so, please elaborate _____

People react differently in stressful situations and may have different coping strategies to deal with the stress of an outbreak. Since the COVID-19 pandemic, have you experienced feelings of:

	1	2	3	4	5
• Uncertainty about the future	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
• Loneliness as a result of social distancing, or self-isolating	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
• Anger or frustration with how the pandemic has disrupted your daily life	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
• Boredom due to being unable to work or engage in regular daily activities/routines	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
• Desire to cope using alcohol or drugs	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
• Sadness, or feelings of hopelessness	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
• Worry or fear about challenges to securing basic needs such as groceries or medication	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

Scale: 1 (least) - 5 (greatest)

Confidential

Page 2 of 3

General Health and Health Management

Do you have a serious or chronic health condition that requires medication and management at home? Yes
 No

If so, please list _____

Has the COVID-19 pandemic affected your daily management of your health conditions? If so, please elaborate. _____

Effects on Housing and Employment

How would you describe your current housing arrangement? I live alone
 I live with spouse/significant other
 I live with another family member
 I live with a non-family member (friend, roommate, etc.)

Did COVID-19 affect your housing situation in any way? (Having someone else move in, having to move out to a new location, limiting the number of people per room...etc.) Yes
 No

If so, please explain _____

Has COVID-19 affected your employment status? Yes
 No

If yes, how so? _____

COVID-19 Exposure and Preventative Measures

In the past 2 weeks, have you experienced any of these symptoms? Check all applicable. Fever
 Dry Cough
 Runny nose
 Body aches
 Upset stomach
 Loss of smell/taste
 Fatigue, tiredness
 None of the above

In the past 3 weeks, have you had close contact with an individual with flu-like symptoms or a confirmed diagnosis of COVID-19? Yes
 No
 Unsure

Have you been tested for COVID-19? Yes
 No

Did you at any point, believe that you might have COVID-19 but DID NOT get tested for any reason? Yes
 No

If yes, please explain _____

Confidential

Page 3 of 3

Was the area you reside in under a "Shelter-in-Place" or "Stay-at-Home" order? Yes
 No
 I don't know

In recent days, have you avoided crowded places whenever possible? Yes
 No

In recent days, have you worn a mask when leaving the house? Yes
 No

What other precautions have you taken to keep yourself and other individuals in your household safe from COVID-19? _____

	1	2	3	4	5
• How worried are you about you or individuals in your household getting COVID-19?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
• How much social distancing do you practice?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
• How often do you practice protective measures such as: hand washing, disinfecting household surfaces, using hand sanitizer?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
• How much control do you feel like you have over whether or not you (or a member of your household) might contract COVID-19?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
• Do you feel like you are equipped with enough knowledge to protect yourself and your household from COVID-19?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

Scale: 1 (least) - 5 (greatest)

How are you typically receiving the latest COVID-19 information and updates?

Check all that apply.

- National news outlet (FOX, CNN, MSNBC, etc.)
- Local news on TV
- Newspaper or local print media
- Directly from family or friends
- A healthcare provider
- Social media (Facebook, Instagram, Twitter, etc.)
- Messaging platform (WhatsApp, Facebook Messenger, etc.)
- Other

If other, please list _____